



## Campylobacteriosis


- Reported cases of disease, 1999-2003
- 5-year rate of disease, 1999-2003
- Reported cases by age group, 2003
- Reported cases by gender, 2003
- Reported cases by county, 2003
- Reported cases by month, 2003



Reported cases of Campylobacteriosis by age group, Rhode Island, 2003			
Age Group	Number of reported cases	Percent	Rate <sup>1</sup> of disease per 100,000
< 1	5	2.8 %	41.0
1-4	12	6.8 %	23.2
5-9	6	3.4 %	8.3
10-19	14	8.0 %	9.5
20-29	19	10.8%	13.9
30-39	25	14.2 %	15.5
40-49	41	23.3 %	25.6
50-59	23	13.1 %	19.8
60-74	19	10.8 %	16.9
75+	12	6.8 %	15.2
<b>TOTAL</b>	<b>176</b>	<b>100.0%</b>	<b>16.8</b>
<sup>1</sup> Denominators for computing rates for RI were based on 2000 Census counts obtained from the U.S. Census Bureau.			

Reported cases of Campylobacteriosis by gender, Rhode Island, 2003			
Gender	Number of reported cases	Percent	Rate <sup>1</sup> of disease per 100,000
Males	87	49.4%	17.3
Females	79	44.9 %	14.5
Missing	10	1.7 %	---
<b>TOTAL</b>	<b>176</b>	<b>100.0%</b>	<b>16.8</b>
<sup>1</sup> Denominators for computing rates for RI were based on 2000 Census counts obtained from the U.S. Census Bureau.			

Reported cases of Campylobacteriosis by county, Rhode Island, 2003			
County	Number of reported cases	Percent	Rate <sup>1</sup> of disease per 100,000
Bristol	7	4.0 %	13.8
Kent	19	10.8 %	11.4
Newport	17	9.7 %	19.9
Providence	101	57.4 %	16.2
Washington	23	13.1 %	18.6
Missing	9	5.1 %	----
<b>TOTAL</b>	<b>176</b>	<b>100.0%</b>	<b>16.8</b>
<sup>1</sup> Denominators for computing rates for RI were based on 2000 population counts obtained from the U.S. Census Bureau.			



Questions regarding these data can be directed to Tara Cooper, MPH: Phone number: (401) 222-3284 or Email: [TaraC@doh.state.ri.us](mailto:TaraC@doh.state.ri.us).